

- Micromelus Silanus, Fem. *Aeneo-niger, cyaneo-varius, antennæ fulvæ, pedes fulvi, alæ fuscæ.*

Corpus æneo-nigrum, subtus cyaneo-nigrum, breve, latum, convexum, parum nitens, scitissime squameum, parce hirtum : caput transversum, breve, thorace latius ; vertex latus ; frons impressa, abrupte declivis : oculi picei, mediocres, non extantes : ocelli vertice triangulum fingentes ; medius perparum antepositus : antennæ fulvæ, ad os insertæ, basi graciles lineares : os piceum : thorax ovatus : prothorax brevissimus : mesothoracis scutum longitudine latius ; parapsides scuto fere in unum confusæ ; paraptera et epimera magna ; scutellum subrhombiforme, postice latius : metathorax brevis, declivis, postice angustus : petiolus brevissimus : abdomen subrotundum, æneum, nitens, læve, supra planum, subtus carinatum, thorace paullo latius et multo brevius : pedes fulvi, simplices, subæquales : alæ fuscæ, mediocres ; squamulæ piceæ ; nervi fusi ; nervus humeralis ulnari duplo longior, radialis ulnari brevior cubitali vix longior, cubitalis sat longus ; stigma mediocre. (Corp. long. lin. $\frac{3}{4}$; alar. lin. $1\frac{1}{4}$.)

Mount Wellington.

VIII.—Descriptions of Chalcidites discovered in St. Vincent's Isle by the Rev. Lansdown Guilding. By FRANCIS WALKER, Esq., F.L.S.

Decatoma Oretilia, Mas et Fem. *Fulvo-flava ferrugineo-varia, antennæ fulvæ, pedes flavi, alæ limpidæ.*

Corpus convexum, sublineare, fulvo-flavum, nitens, subtilissime punctatum, parce hirtum : caput transversum, breve, thorace paullo latius ; vertex latus ; frons impressa, abrupte declivis : oculi rufi, mediocres, non extantes : antennæ fulvæ, clavatae, thorace non longiores ; articulus 1^{us} longus, gracilis ; 2^{us} cyathiformis ; 3^{us} et 4^{us} minimi ; 5^{us} et sequentes breves, usque ad 9^{um} latescentes ; clava fusiformis, acuminata, articulo 9° multo latior et plus duplo longior : thorax ovatus : prothorax maximus, transversus, subquadratus, postice ferrugineo-maculatus : mesothoracis scutum brevissimum ; parapsidum suturæ vix conspicuæ ; scutellum subrotundum ; discus ferrugineus : metathorax transversus, brevis : petiolus brevis : abdomen ovatum, læve, mari thoracis longitudine, fem. thorace paullo longius subtus carinatum apice acuminatum fuscum ; vitta per dorsum interrupta fusca : pedes pallide flavi, simplices, subæquales : alæ limpidæ ; squamulæ fulvæ ; nervi fulvi ; nervus humeralis ulnari fere duplo longior, radialis ulnari non brevior cubitali plus duplo longior ; stigma minutum. (Corp. long. lin. 1 ; alar. lin. $1\frac{1}{2}$.)

Pteromalus Helice, Fem. *Aeneo-viridis, antennæ fulvæ, pedes flavi, alæ limpidæ.*

Corpus convexum, æneo-viride, nitens, scitissime squameum, parce hirtum : caput transversum, breve, thorace latius ; vertex latus ; frons impressa, abrupte declivis : oculi rufi, mediocres, non extantes : antennæ fulvæ, extrorsum crassiores, thorace vix longiores ; articulus 1^{us} longus, gracilis ; 2^{us} cyathiformis ; 3^{us} et 4^{us} minimi ; 5^{us} et sequentes breves, usque ad 10^{um} latescentes ; clava longiconica, acuminata, articulo 10° duplo longior : thorax ovatus : prothorax transversus, brevissimus : mesothoracis scutum longitudine latius ; parapsidum suturæ vix conspicuæ ; scutellum subconicun, ante-truncatum : metathorax obconicus, mediocris, declivis : petiolus brevissimus : abdomen fusiforme, læve, supra depresso, subtus carinatum, apice acuminatum, thorace longius et angustius : pedes læte flavi, simplices, subæquales : alæ limpidæ ; squamulæ fulvæ ; nervi flavi ; nervus humeralis

ulnari multo longior, radialis ulnari vix brevior, cubitalis sat longus; stigma minutum. (Corp. long. lin. 1— $1\frac{1}{4}$; alar. lin. $1\frac{3}{4}$ —2.)

Var. β. Thorax æneus.

Genus LELAPS, *Haliday MSS.*

Fem. Corpus angustum, sublineare, convexum, nitens, scitissime squameum, subtus lœve, parce hirtum: caput transversum, breve, thoracis latitudine; vertex angustus; frons impressa, abrupte declivis: oculi magni, non extantes: ocelli vertice triangulum fingentes: antennæ gracillimæ, extrorsum crassiores, pubescentes, corpore paullo breviores; articulus 1^{us} longus, gracilis; 2^{us} longicyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes longi, usque ad 10^{um} curtantes et latescentes; clava linearis, apice obtusa, articulo 10^o longior: thorax longiovatus: prothorax brevis, antice angustus: mesothoracis scutum longitudine latius; parapsidum suturæ bene determinatæ, postice approximatæ; scutellum obconicum?: metathorax mediocris, obconicus, declivis: petiolus brevissimus: abdomen longiovatum, breve, subtus profunde carinatum, thorace longius; segmentum 1^{um} maximum: oviductus exsertus; vaginalæ abdomine paullo breviores: pedes longi, graciles, simplices, subæquales; tarsorum articuli 1^o ad 4^{um} curtantes, 5^{us} 4^o paullo longior; unguis et pulvilli minimi: alæ mediocres; nervus humeralis ulnari non longior, radialis ulnari brevior alæ apicem fere attingens, cubitalis sat longus; stigma minimum.

Lelaps pulchricornis (*Haliday MSS.*), *Fem.* *Cuprea, antennæ fuscæ flavocinctæ, pedes flavi, proalæ fusco-nebulosæ.*

Corpus cupreum, viridi-varium: oculi et ocelli rufi: antennæ fuscæ, flavocinctæ, apice flavæ; articulus 1^{us} flavus: oviductus vaginalæ fulvæ, apice fusca: pedes flavi; unguis et pulvilli fusi: alæ limpidæ; proalæ fusco-nebulosæ; squamulæ fulvæ; nervi fulvi. (Corp. long. lin. 1 $\frac{1}{2}$; alar. lin. 2 $\frac{3}{4}$.)

Genus IDARNES.

Corpus sublineare, depresso, lœve, nitens, parce hirtum: caput transversum, breve, thoracis latitudine; vertex sat latus: palpi biarticulati: oculi mediocres, non extantes: antennæ breves, clavatae; articulus 1^{us} longus, validus; 2^{us} cyathiformis; 3^{us} et sequentes brevissimi: thorax ovatus: prothorax magnus, transversus: mesothoracis scutum impressum, longitudine latius; parapsidum suturæ remotæ, bene determinatæ, postice approximatæ; scutellum magnum, transversum, bisulcatum: metathorax magnus, subquadratus: petiolus vix ullus: abdomen longi-ovatum, thorace paullo longius non latius; segmentum 1^{um} magnum, 2^{um} et sequentia brevia: oviductus longissimus; vaginalæ corpore triplo longiores: propedum coxae magnæ; femora crassa; tibiæ latæ: mesopedes graciles; coxæ parvæ: metapedum coxae sat magnæ; femora crassa; tibiæ subarcuatæ: tarsorum articuli 1^o ad 4^{um} curtantes, 5^{us} 4^o paullo longior; unguis et pulvilli minimi: alæ mediocres; nervus humeralis longus, ulnaris brevissimus, radialis longus ulnari vix brevior, cubitalis longissimus, arcuatus, ulnari longior; stigma minimum.

Idarnes Carme, Fem. *Viridi-æneus, abdomen cupreum, antennæ nigræ, pedes fulvi, alæ limpidæ.*

Corpus viridi-æneum: oculi et ocelli rufi: antennæ nigræ: abdomen cupreum: oviductus fulvus; vaginalæ nigræ: pedes fulvi; unguis et pulvilli fusi: alæ limpidæ; squamulæ fulvæ; nervi flavi. (Corp. long. lin. 1; alar. lin. 1 $\frac{3}{4}$.)

Encyrtus Gargaris, Mas. *Viridis, abdomen cupreum, antennæ fulvae, pedes fulvi, alæ albo-limpidæ.*

Corpus breve, convexum, viride, nitens, scitissime squameum, parce

hirtum : caput transversum, breve, thoracis latitudine ; vertex latus ; frons impressa, abrupte declivis : oculi picei, mediocres, non extantes : antennæ fulvæ, submoniliformes, extrorsum crassiores, thorace non longiores ; articulus 1^{us} longus, gracilis ; 2^{us} cyathiformis ; 3^{us} et sequentes transversi ; clava conica : thorax ovatus : prothorax minimus, supra non conspicuus : mesothoracis scutum magnum, planum, longitudine non latius ; parapsidum suturæ non conspicuæ ; scutellum obconicum : metathorax transversus, brevissimus : petiolus vix ullus : abdomen depresso, obconicum, cupreum, lœve, thorace multo brevius : pedes fulvi ; mesotarsi et metatarsi pallidiores, apice fuscæ : alæ albo-limpidae ; squamulae piceæ ; nervi fulvi ; nervus humeralis nullus, ulnaris costæ dimidiæ longitudine, radialis brevissimus, cubitalis sat longus ; stigma minutum. (Corp. long. lin. $\frac{1}{2}$; alar. lin. $\frac{3}{4}$.)

Reared from a larva.

Euplectrus Furnius (bicolor?), Mas. *Ater, abdominalis discus fulvus, antennæ fuscæ, pedes fulvi, alæ limpidae.*

E. bicolori simillimus. Corpus convexum, atrum, nitens, lœve, pilis albis parce hirtum : caput transversum, brevissimum, impressum, vix thoracis latitudine ; vertex latus ; frons abrupte declivis : oculi rufi, mediocres, vix extantes : antennæ fuscæ, filiformes ; thorace longiores ; articulus 1^{us} longus, gracilis, flavus, apice fuscus ; 2^{us} cyathiformis ; 3^{us} et sequentes ad 6^{um} mediocres, subæquales ; clava ovata, articulo 6° longior : thorax ovatus : prothorax angustus, mediocris, conicus : mesothoracis scutum longitudine latius ; parapsidum suturæ non bene determinatæ ; scutellum subovatum : metathorax obconicus, sat magnus, declivis : petiolus gracilis, sat longus : abdomen subconicum, depresso, thorace multo brevius ; segmentum 1^{um} magnum, 2^{um} et sequentia brevia ; discus fulvus : pedes fulvi, simplices, subæquales ; unguis et pulvilli fuscæ ; mesotibiae apice spina sat longa armatae : alæ limpidae ; squamulae fulvæ ; nervi flavi ; nervus humeralis ulnari multo longior, radialis ulnari multo brevior cubitali paullo longior ; stigma minutum. (Corp. long. lin. 1 ; alar. lin. 1 $\frac{3}{4}$.)

PAPHAGUS, n. g.

Fem. Corpus sublineare, depresso, nitens, lœve, parce hirtum : caput transversum, breve, thoracis latitudine ; vertex latus ; frons impressa : oculi mediocres, non extantes : antennæ clavatae, pubescentes, prope os insertæ, thorace non longiores ; articulus 1^{us} validus, sublinearis ; 2^{us} longicyathiformis ; 3^{us} et sequentes usque ad 6^{um} latescentes ; clava ovata, acuminata, articulo 6° longior et latior : thorax ovatus : prothorax transversus, brevis : mesothoracis scutum longitudine latius ; parapsidum suturæ non bene determinatae ; scutellum transversum, latum : metathorax mediocris, transversus, postice angustus : petiolus brevis : abdomen ovatum, supra depresso, subtus carinatum, apice acuminatum, thorace non longius : pedes simplices, graciles, subæquales ; tarsorum articuli 1° ad 4^{um} curtantes ; unguis et pulvilli minimi : alæ mediocres ; nervus ulnaris humerali paullo longior, radialis nullus, cubitalis brevis ; stigma minutum.

Paphagus Sidero, Fem. *Cupreus, abdomen basi cyaneum, antennæ fuscæ, pedes flavi, alæ limpidae.*

Cupreus : oculi picei : antennæ fuscæ ; articulus 1^{us} flavus : mesothoracis scutum viridi-varium ; scutellum purpureo-varium : metathorax viridi-cyanus : petiolus fulvus : abdomen basi cyaneum : pedes flavi ; unguis et pulvilli fuscæ : alæ limpidae ; squamulae fulvæ ; nervi flavi. (Corp. long. lin. 1 ; alar. lin. 1 $\frac{1}{2}$.)

Telenomus Apilius, Fem. *Ater, antennæ nigrae, pedes picei, tarsi fulvi, alæ albo-limpidae.*

Corpus breve, latum, crassum, convexum, atrum, rude rugulosum, parum

nitens, parce hirtum : caput transversum, breve, thoracis latitudine; vertex latus; frons convexa, abrupte declivis: oculi picci, mediocres, non extantes: antennæ clavatæ, nigræ, corpore breviores; articulus 1^{us} sublinearis, validus; 2^{us} longicyathiformis; 3^{us} et 4^{us} angusti; 5^{us} et sequentes latiores, clavam fingentes longam fusiformem: thorax ovatus: prothorax brevissimus, supra non conspicuus: mesothoracis scutum magnum, longitudine latius; parapsidum sutura non conspicua; scutellum mediocre, brevi obconicum: metathorax transversus, brevissimus: petiolus vix ullus: abdomen subovatum, scite rugulosum, thorace non longius; segmenta basi sulcata, 1^{um} magnum, 2^{um} et sequentia brevia: pedes picci, simplices, subæquales; trochanteres fulvi; genua fulva; tibiae basi fulvae; tarsi fulvi, apice fusi: alæ albolimpidae; squamulæ fuscæ; nervi fulvi; nervus humeralis nullus, ulnaris costæ dimidio brevior, cubitalis longus radiali paullo brevior; stigma minimum. (Corp. long. lin. 2; alar. lin. 1.)

Reared from the eggs of an insect.

BIBLIOGRAPHICAL NOTICES.

Manual of British Botany, containing the Flowering Plants and Ferns, arranged according to the Natural Orders. By Chas. C. Babington, M.A., F.I.S., F.G.S., &c. &c. London: J. Van Voorst, 1843.

THE present work is modelled after the well-known ‘Synopsis Floræ Germanicæ’ of the learned Koch, and is the result of a critical examination of our native plants, by a careful comparison of indigenous specimens with the works of eminent continental authors and with plants obtained from other parts of Europe, whereby a desideratum long wanted has been attained, viz. the assimilation of the nomenclature of British and continental writers. To effect this, our flora has undergone a general and scrutinizing revision, so as to determine what plants are really deserving of the rank of distinct species, and what must be regarded only as varieties; and those names have been adopted which possess the claims of priority by a careful examination of all the best European floras, comparing our plants with the descriptions contained in them, and in very many cases with foreign specimens of undoubted authenticity. In the adoption of genera and species an endeavour has been made, by the examination of the plants themselves, to determine which are truly distinct, thus taking nature as a guide, and not depending upon the authority of any name, however distinguished. As the volume is intended as a field-book, or travelling companion for botanists, the characters have been admirably and ably contrasted, and condensed as much as possible consistent with their accurate discrimination. Synonyms have been wholly omitted, except quotations of one British and one German figure to each plant. Localities are only given for new or peculiarly rare plants, though the general distribution through the British Isles, or in the several main portions thereof, and their character in that distribution, have been noted. The arrangement is that of the Natural System, but to render it generally available, a succinct synopsis of the Linnaean genera is added. To detail its many excellences would exceed our due limits, for fully to appreciate the scientific and critical accuracy and discrimination with which the